

ACT!ON



Advanced multitemporal modelling and optimisation of CO₂ Transport, stOrage and utilisation Networks

WP 5. Task 5.5. - ROMANIAN CLUSTER

November 2023


STATUS OF CCUS IN ROMANIA



Work related to the CO₂ geological storage began in Romania with the affiliation of the GeoEcoMar to ENeRG in 2002 and continued with participation of the institute in international and national projects related to CCS:



- subcontractor in "CASTOR" project,
- partner in "EU GeoCapacity", "CO₂ Net East", "Impact of communication", "CGS Europe", "CO₂ Stop" European projects,
- partner in similar national projects: "The National Program of Carbon Capture and Storage for 2011-2020 period" and "Geological storage" section of the Feasibility Study for the "Getica CCS" Demonstration Project.

After accession to the CO₂ GeoNet Association on 2013, GeoEcoMar is participating in the "ENOS" HORIZON 2020 project for 2016-2020 period. Since 2017 GeoEcoMar was partner in ACT projects: "ALIGN CCUS", "ECO-BASE", " REX-CO₂", as well as, since 2022, in ACTION project.



CO₂ Major Emissions and the Geological Storage Capacity in Romania


GETICA CCS DEMONSTRATION PROJECT




2 zones have been selected as best candidates for storage

- Approx 2000 m depth
- Reservoir: Sarmatian sandstone formation from Getic Depression
- Caprock: Upper Sarmatian shaly formation
- Static and dynamic modelling


- Capture: Turceni powerplant
- Transport: onshore pipelines
- Storage: 1.5 Mt of CO₂ to be stored in deep saline aquifers
- Planned to start injection at the end of 2015
- Project competed for NER 300, high scored, on the waiting list
- Loss of governmental support lead to stalling the project



Zone 1



Zone 5






MULTIMODAL TRANSPORT OF CO₂

The development of a CO₂ transport infrastructure is essential for speeding up the implementation of CO₂ capture and storage technologies in Romania. This infrastructure does not necessarily rely on onshore pipelines only depending on the height of the capital costs for developing such an infrastructure versus the investments in ship transport.

The best solution would be to combine transport by pipeline with transport by vessels. This is also an option for Romania, taking into account the existence of the Danube, of the Danube - Black Sea Canal and the opening to the Black Sea where good storage reservoirs have been identified.


So, multimodal transport of CO₂ consists in a smart usage of pipelines and ships. Instead of a unique network of pipelines, multimodal transport of CO₂ means a large usage of specialized ships on the inland waterways, and short pipelines between the emission sources as well as suitable storage locations with the closest harbours.




National road network



National railways network



Romanian Oil Pipelines



Romanian Gas Pipelines

Originating in Germany, the Danube flows southeast for 2,850 km, passing through or bordering:
- Austria, Slovakia, Hungary, Croatia, Serbia, Romania, Bulgaria, Moldova, and Ukraine.






TEN-T road network and inland waterways on Romania and some of neighbouring countries



Romanian Danubian and maritime Ports



Map of ports on the Danube river, major CO₂ emissions on Southern part of Romania and hydrocarbon deposits suitable for EOR


ROMANIAN CCUS CLUSTER

From the analysis of major CO₂ emission sources in the southern part of the country, we can conclude that these could be grouped into a major cluster for the purpose of CO₂ transport.

In our vision, the best solution would be to design small segments of pipelines for the CO₂ transport from an individual source to a central or nodal point at which all the captured emissions are collected. The next segment of transport is also a pipeline, from the nodal point to a Danube harbour, at which the CO₂ will be transferred to a barge or vessel and transported by river to the offshore region where it will be injected for CO₂ EOR operations or for storage in deep saline aquifers.

CCUS stands for Carbon Capture, Utilisation, and Storage. A CCUS cluster typically involves the integration of various components:

- carbon capture facilities,
- transportation infrastructure for the captured CO₂,
- utilisation facilities where the CO₂ is put to use in industrial processes (such as in industrial processes or for enhanced oil recovery),
- storage sites for the secure and long-term underground storage of CO₂.




Oltenia CCUS Cluster

The clustering of these elements allows for a more efficient and coordinated approach to carbon management, contributing to the reduction of greenhouse gas emissions and addressing climate change challenges.

Also, a CCUS cluster involves collaboration between different industries, government bodies, and research institutions to develop and deploy these technologies on a larger scale, aiming to reduce greenhouse gas emissions and mitigate climate change.

The clustering of these activities allows for shared infrastructure, knowledge, and resources, making the overall implementation of CCUS more efficient and effective.




CO₂ EOR clusters



The blueprint for the Romanian Cluster refers to:

- identification of the most feasible transportation routes for future captured CO₂,
- investigation of the geological storage solutions available, including the possibility to use the CO₂ for enhanced hydrocarbon recovery in the region.

The existence of Danube at the southern border of Romania presents a very good opportunity for the transport of the captured CO₂ emissions toward the Black Sea.




Major CO₂ Emissions and Hydrocarbon Deposits in Romania



- Good connection for the southern part of Romania due to the proximity to the Danube
- Good connection to the Black Sea and its storage and EOR possibilities in the Histria Depression area

- Opportunities in Histria Depression (Black Sea) - an important unit for hydrocarbon exploration and exploitation.
- Potential storage sites in deep saline aquifers *Iris*, *Lotus* and *Tomis*, located on the southern flank of the Histria Depression
- good reservoir formations in Albian, Lower and Upper Cretaceous and Middle Eocene. Protected by shale sequences on top and secondary by the Oligocene shale formation.



ACKNOWLEDGEMENTS

This project is funded through the ACT - Accelerating CCS Technologies programme. Financial contributions by the Department for Business, Energy & Industrial Strategy UK (BEIS); the Ministry of Economic Affairs and Climate Policy, the Netherlands; Department of Energy-Office of Fossil Energy (DOE-FE); Emissions Reduction Alberta (ERA); the French Agency for the Environment and Energy Management (ADEME); **the Executive Agency for Higher Education, Research, Development and Innovation Funding (UEFISCDI)** are gratefully acknowledged. We also thank all our industrial partners for their contributions and the funding they provide towards ACTION.

Website: www.action-act.org

Twitter: @ACTION_ACT

THANK YOU FOR YOUR ATTENTION!