

VOLHYNIAN MICROFAUNA IN BLEJEȘTI AREA (CENTRAL-SOUTHERN PART OF THE MOESIAN PLATFORM)

CORINA IONIȚĂ-BADEA^{1,2}, OVIDIU DRAGASTAN², MIHAELA C. MELINTE-DOBRINESCU³,
ANDREI BRICEAG³, COSTEL-VICTOR SINGHEL¹

¹OMV Petrom, ICPT Laboratory Câmpina, 29 Culturii Boulevard, Câmpina, Prahova County,
Romaniacorina.ionita@petrom.com

²University of Bucharest, Faculty of Geology and Geophysics, Department of Geology, Bălcescu Boulevard 1, 010041

³National Institute of Marine Geology and Geo-Ecology (GeoEcoMar), 23-25 Dimitrie Onciul St., 024053 Bucharest, Romania

Abstract. This paper presents the microfaunal assemblages of lower Sarmatian (Volhynian) deposits intercepted by 10 hydrocarbon wells from the Blejești hydrocarbon field, S Romania. The identified microfaunas are typically for two Volhynian foraminifera zones: *Varidentella reussi* Zone and *Elphidium reginum* Zone. At the contact with the underlying Cretaceous or Middle Miocene (= Badenian) units, the Sarmatian studied samples, containing foraminiferal assemblages with abundant *Ammonia beccarii*, may not be assigned to a certain biozone. Besides, the identified ostracod assemblages were assigned to the NO 11 *Cytheridea hungarica* – *Aurila mehesi*. The composition and the abundance of microfaunal assemblages allow us to advance paleoecological considerations for the lower Sarmatian deposition interval of the studied area.

Key words: early Sarmatian, foraminifera, ostracods, biozones, paleoecology, S Romania

1. INTRODUCTION

The studied area is located N of the Danube river and belongs to the central-southern region of the Moesian Platform, tectonically framed in the Roșiori - Alexandria Depression. The Moesian Platform (Fig. 1) is a major tectonic unit, bordered in the N by the Getic Depression, the Central Dobrogea towards NE, the Balkan Mountains in the S and the Black Sea in the E (Săndulescu, 1984).

The Moesian Platform is divided (Săndulescu, 1984) in: (i) the western part, namely Wallachian platform that extends up to the Danube; (ii) the eastern part, *i.e.*, the Southern Dobrogea platform, bordered by the Danube to the West and the Black Sea towards East. The Wallachian part of the Moesian Platform contains many hydrocarbon wells that still produce nowadays and therefore it was intensively studied since the beginning of the Romanian oil industry, due to its high potential (Paraschiv, 1979; Beca & Prodan, 1983).

Within the Miocene, the western Moesian Platform was included in the Dacian Basin, part of the Central Paratethys. Since the Upper Miocene (*i.e.*, Sarmatian stage)

the Dacian Basin was included in the Eastern Paratethys, due to paleogeographic changes (Popov *et al.*, 2004; Jipa & Olariu, 2009).

The Paratethyan Domain occurs since the Oligocene, as a result of the tectonic movements Alps and Carpathian, following the African-Arabian Plate collision with the Eurasian one (Royden *et al.*, 1982; Steininger & Wessely, 2000). Starting with Middle-Late Miocene the Paratethys was divided in several domains grouped in Central Paratethys, that included Vienna, Pannonian and Transylvanian basins and the Eastern Paratethys with Dacian, Euxinian and Caspian basins (Papp *et al.*, 1974; Marinescu, 1978; Popov *et al.*, 2004; Piller *et al.*, 2007) (Fig. 2). Since the Late Miocene and later, in the Pliocene-Quaternary interval, the original Oligocene extension of the Paratethys Sea significantly decreased, and isolated basins, *i.e.*, including the present day Black Sea, along with the Caspian and Aral ones, occurred. Therefore, the paleogeographic changes mirrored the significant uplift of the Carpathians and Caucasus mountains (Popov *et al.*, 2004; Schmid *et al.*, 2008).




Fig. 1. Geological map of Romania (simplified after Săndulescu, 1984) showing the location of the investigated wells.




Fig. 2. Paleogeography of the Paratethyan Domain and surrounding areas during the Sarmatian (modified after Popov *et al.*, 2004, Jipa & Olariu, 2009 and Briceag *et al.*, 2018).

This paper is focused on the Middle Miocene, *i.e.*, Sarmatian *sensu lato* (*s.l.*) stage, from wells of the W Moesian Platform. The Sarmatian regional stage was named by Suess (1866) and supported over time many discussions regarding its chronostratigraphic division. In the Central Paratethys, the Sarmatian *sensu stricto* (*s.s.*) proposed by Suess (1866) is in used (Piller *et al.*, 2007); there, the Sarmatian stage (containing two substages, the Volhynian and the Bessarabian) overlies the Badenian stage and it is followed by the Pannonian; in the Eastern Paratethyan Domain, the Sarmatian *s.l.*, comprised between the Badenian and Meotian stages, is used. There, the Sarmatian is divided into three substages (Fig. 3), namely Volhynian, Bessarabian and Khersonian (Andrusov, 1917; Marinescu, 1978; Popov *et al.*, 2004; Brănzilă & Chira, 2005; Jipa & Olariu, 2009, 2013; Briceag *et al.*, 2018).

2. MATERIAL AND METHODS

Qualitative and quantitative foraminiferal and ostracod analysis have been performed. In all, 73 samples have been collected from cores belonging to 10 wells from the Blejești hydrocarbon field, situated between 60 and 80 km SE from Bucharest, in the Teleorman County (Fig. 4). Each collected sample weighs about 100 grams. For microfaunal analysis, the material was processed by standard micropaleontological methods: the material was soaked in water containing 30% hydrogen peroxide, boiled and passed through 63 µm sieve. All the specimens have been examined under a binocular Olympus microscope and microphotographs have been taken with an Olympus camera. The most common encountered taxa are illustrated in Plates 1-4.

To assign biostratigraphically the identified assemblages, we used the Central Paratethys biozonation (Grill, 1941; Papp, 1974, Rögl, 1998; Görög, 1992) along with the biozonation schemes defined in various regions of Romania, *i.e.*, Ionesi, 1968; Popescu, 1995; Filipescu & Silye, 2008 (Fig. 3). All the data, including samples, thickness of Volhynian formations intercepted by wells, age, number of taxa along with the lithology have been used for making graphics. To note that the figure given to each of the studied well has been attributed by the authors, does not match with the real number of the wells.

3. RESULTS AND DISCUSSION

In the studied wells, the stratigraphic interval assigned to the early Volhynian reaches almost 100 m in thickness. Predominantly type of sediments are: medium to low compacted argillaceous limestones with horizontal laminae (in some places commonly containing pyrite and organic matter), argillaceous marls with carbonized plant remains, sands (fine grained, subangular to subrounded, moderately to well sorted), calcareous marls (medium compacted, fine micaceous), sandstones (fine-medium grained, coarse at places, subangular to subrounded, moderately sorted, low to medium cemented with mixed binder, fine micaceous) and slightly argillaceous limestones with oolites (well compacted and containing fossil debris).

The Volhynian deposits overlies the Badenian sediments (only in Well Blejești 3) or the Cretaceous ones (in 6 wells); the base of the Volhynian contains very scarce microfaunal assemblages. In the claystones of Well Blejești 3, above the

SARMATIAN FORAMINIFERA BIOZONES IN PARATETHYS																
Austria		Czech Republic Slovakia		Hungary		Romania (Transilvanyan Basin), Popescu, 1995; Filipescu and Syle, 2008		Romania (Moldavian Platform) Ionesi, 1968								
Grill, 1941; Papp, 1974; Rögl, 1988		Jiríček, 1972		Boda, 1974; Görög , 1992												
Pannonian s. str.						Pannonian		Maeotian								
						Bessarabian Substage		Porosonion aragviensis/Streptochilus		Khersonian						
						Dogielina sarmatica		Nonion bogdanowiczii, Porosonion subgranosus, Elphidium macellum								
								Porosonion subgranosus and Elphidium macellum								
Nonion granosum		Late Sarmatian	Porosonion subgranosum	Porosonion subgranosum	Bessarabian Substage		SARMATIAN s. l.									
				Spirolina austriaca												
		Early Sarmatian	Elphidium hauerinum	Elphidium hauerinum							Volhynian substage	Elphidium reginum	Articulina sarmatica			
														Elphidium reginum	Rotalia beccarii, Porosonion subgranosus,	Peneroplidae
															Elphidium rugosum	Rotalia beccarii and Quinqueloculina consobrina Subzone
	Rotalia beccarii Subzone															
Elphidium reginum		Elphidium reginum				Varidentella reussi		Quinqueloculina reussi, Articulina problema								
								Zone with rare foraminifera taxa (Elphidium, Nonion, Globigerina)								
Anomalinoides divdens		Cibicides badenensis	Elphidium reginum					Elphidium regina								
								Cibicides lobatulus, Quinqueloculina fluviata, Q. karrerovata								

Fig. 3. Central and Eastern Paratethys Sarmatian substages and correlation based on foraminifers.



Fig. 4. The location of the Blejești studied wells (map from www.google.com/maps)

Badenian deposits, the argillaceous limestones are very abundant in macrofossils, while the microfaunas are missing or are inconclusive: sponge spicules and rare *Charophyta* algae fragments along with the benthic foraminifera species *Ammonia beccarii* (Linnaeus). Similar fossil assemblages collected from the deposits overlaying the top of the Badenian were described in a nearby area, in wells belonging to the Cartojani oil field (Ioniță *et al.*, 2016).

Above this basal level, the rest of the cores, assigned to the early Volhynian, contains very fine polymictic sands with scarce microfaunal assemblages, composed of sponge spicules, few benthic foraminifera, such as *Elphidium rugosum* (d'Orbigny), *Elphidium aculeatum* (d'Orbigny), *Nonion bogdanowiczi* Voloshinova, rare ostracods specimens of *Cytheridea hungarica* (Zalányi), *Charophyta* algae fragments and reworked Cretaceous foraminifera (many of them with pyritized test). We assume that a brackish paleoenvironment prevailed within the lower part of the Volhynian, based on the aforementioned benthic foraminifers.

The overlaying deposits contain microfaunas with very abundant *Ammonia beccarii*, between 100 – 1400 taxa/sample). The tests of this species are small-sized and most of them are filled with pyrite (Fig. 5), suggesting low oxygenation in the marine bottom water. This assemblage is completed by few specimens of benthic foraminifers *Elphidium macellum* (Fichtel & Moll) and *Cycloforina* sp., along with ostracods such as *Aurila mehesi* (Zalányi), *Cytheridea hungarica* and *Miocyprideis sarmatica* (Zalányi); rarely, mysidae statoliths occur. Besides *in situ* taxa aforementioned, Cretaceous and Miocene reworked foraminifer species were observed.

The above-mentioned ostracod assemblage argues for the presence of the Central Paratethys *Cytheridea hungarica* - *Aurila mehesi* ostracod zone (Fig. 6), early Sarmatian in age (Jiříček & Říha, 1991; Filipescu *et al.*, 2014). This biozone is described also from extra Carpathian regions, *i.e.*, the Dacian Basin extending on the Moldavian Platform (Ionesi & Guevara, 1993; Dumitriu *et al.*, 2017), occurring in fact in the lower Sarmatian depositional interval around the whole Paratethyan Domain (Zelenka, 1990).

The foraminifer *Ammonia beccarii*, commonly present in the studied samples, is an euryhaline species, adapted to high salinity fluctuations (Koubová & Hudáčková, 2010). Some authors (Gross, 2004; ter Borgh *et al.*, 2014) indicate that the ostracod species *Cytheridea hungarica* is indicative for an epineritic, brackish to normal marine setting. Other authors suppose that this taxon is more related to a shallow littoral to sublittoral settings, requiring well aerated brackish waters, with depth no more than 30 m and a warm climate, *i.e.*, tropical to subtropical (Jiříček, 1985; Olteanu & Jipa, 2006). Assemblages with dominant *Elphidium* spp. and *Ammonia beccarii* are currently present in brackish waters all over the world, at salinity above 4-5 ‰ (Murray, 1991). According to Reinhardt *et al.* (1994), the domination of *Ammonia* is linked, in terms of paleosetting, to a transitional zone between lagoon and open marine conditions. Debenay *et al.* (1998) indicate that *Ammonia beccarii* is common in marine environments with salinities greater than 33‰. On the other hand, *Ammonia beccarii* is present nowadays in the Black Sea, including near the Danube mouth, at salinities around 5-8 ‰ and in some Pleistocene-Holocene intervals known to have had salinity strong fluctuations (Melinte-Dobrinescu & Ion, 2013; Briceag *et al.*, 2016a, 2016b, 2019).




Plate 1. Volhynian foraminifera from Blejești area. **1.** *Cycloforina* sp. – side view, **2.** *Vatricentella reussi* Bogdanowicz – side view, **3.** *Articulina problema* Bogdanowicz – side view, **4.** *Elphidium aculeatum* (d'Orbigny) – spiral view, **5.** *Elphidium crispum* (d'Orbigny) – spiral view, **6.** *Elphidium fichtelianum* (Linnaeus) – spiral view, **7.** *Elphidium hauerinum* (d'Orbigny) – spiral view, **8.** *Elphidium josephinum* (d'Orbigny) – spiral view, **9.** *Elphidium macellum macellum* (Fichtel & Moll) – spiral view.




Plate 2. Volhynian foraminifera from Blejești area. **10.** *Elphidium macellum aculeatum* (Silvestri) – spiral view, **11.** *Elphidium reginum* (d'Orbigny) – Blejești 3 well, spiral view, **12.** *Elphidium reginum* (d'Orbigny) – Blejești 2 well, spiral view, **13.** *Lobatula lobatula* (Walker and Jakob) – umbilical view (left) and spiral view (right), **14.** *Nonion bogdanowiczi* Voloshinova – spiral view, **15.** *Ammonia beccarii* (Linnaeus) – umbilical view (left) and spiral view (right), **16.** *Fissurina* sp. – side view.




Plate 3. Völnynian ostracods from Blejești area. All valves are in external lateral view. LV = left valve, RV = right valve. **1.** *Cytheridea hungarica* (Zalányi) – RV, **2.** *Aurila mehesi* (Zalányi) – with pyritized, LV, **3.** *Aurila mehesi* (Zalányi) – LV, **4.** *Aurila merita* (Zalányi) – LV, **5.** *Amnicythère tenuis* (Reuss) – RV, **6.** *Amnicythère tenuis* (Reuss) – infilled with pyrite, RV.



Plate 4. Volhynian ostracods from Blejești area. All valves are in external lateral view. LV = left valve, RV = right valve. **7.** *Callistocythere incostata* Pietrzeniuk – RV, **8.** *Callistocythere egregia* (Méhés) – LV, **9.** *Euxinocythere diafana* (Stancheva) – LV, **10.** *Cytherois sarmatica* (Jiříček) – RV, **11.** *Loxoconcha curiosa* Schneider – LV, **12.** *Loxocorniculum schmidi* (Cernajsek) – LV, **13.** *Xestoleberis elongata* Schneider – RV, **14.** *Xestoleberis fuscata* Schneider – RV.

The foraminifera species *Ammonia beccarii* is seen by many authors as an opportunistic taxon. Hence, its dominance in the assemblages is most probably indicative for a low degree or even absence of competition with other taxa (Murray, 1991).




Fig. 5. Numerous specimens of *Ammonia beccarii* from Blejești 3 associated with few *Cycloforina* fragments and *Miocyprideis sarmatica* ostracod species.

In the identified ostracod assemblage of the studied samples, *Cytheridea hungarica* is the dominant ostracod species. Its common presence, along with the significant abundance of foraminifer *Ammonia beccarii*, constituting monospecific assemblages, indicates a brackish to shallow marine paleoenvironment, located close to the shoreline.

This assumption is also supported by the presence of *Charophyta* algal fragments, possible transported by rivers from the adjacent land. Along with the aforementioned assemblages, gastropods, bivalve fragments, rare fish bones and worm tubes suspension feeder *Serpula* are present. The common presence of *Serpula* taxa in the various

Paratethyan Sarmatian deposits was interpreted to be linked to a progressive basin desalting and hence the dominance of more brackish conditions, but a marine shallow paleosetting is not excluded (Saint-Martin & Pestrea, 1999; Cornée *et al.*, 2009). Nowadays, the presence of abundant *Serpula* shows an enclosed and sheltered environment with no turbulent water, mainly in a marine setting, frequently associated with reefs (Short & Neckles, 1999).

The overlaying sediments are represented by very fine polymictic sands with scarce microfaunal assemblages, including sponge spicules and few benthic foraminifera taxa: *Elphidium rugosum* (d'Orbigny), *Elphidium aculeatum* (d'Orbigny) and *Nonion bogdanowiczii* Voloshinova, rare ostracods belonging to the species *Cytheridea hungarica* (Zalányi), *Charophyta* algae fragments and various reworked foraminifera, many of them with pyritized test (Plates 1-4). The above-described microfaunal assemblages are most probably related to a brackish paleosetting.

The overlaying deposits of the studied cores contain foraminiferal assemblages characteristic for *Varidentella reussi* Zone that includes many miliolid taxa, such as *Cycloforina fluviata* Venglinskin, *Quinqueloculina akneriana akneriana* (d'Orbigny), *Varidentella reussi* (Bogdanowicz), *Pseudotriloculina consobrina* (d'Orbigny) and ostracods, i.e., *Aurila mehesi* (Zalányi), *Aurila merita* (Zalányi), *Callistocythere egregia* (Méhes), *Loxoconcha curiosa* Schneider and *Xestoleberis fuscata* Schneider. Similar microfaunal association was mentioned from boreholes of the Moldavian Platform – NE Romania (Dimitriu *et al.*, 2017).

The *Varidentella reussi* Foraminiferal Zone extends from immediately above (5 m stratigraphically) the association with abundant *Ammonia beccarii*. Because of the foraminifer index species absence in the studied wells, the oldest Sarmatian biozone *Anomalinoidea dividens* Zone (Papp *et al.*, 1974; Popescu, 1995; Hartzhauser & Piller, 2004; Filipescu & Silye, 2008) was not identified, thus we may assume that the lower Volhynian depositional interval is lacking due to the erosion or nondeposition.

SARMATIAN OSTRACODES ZONES FROM PARATETHYS (Slovakia, Czech Republic, Bulgaria)				
AGE	ZONE	Jiríček & Riha, 1991	Zelenka, 1990	Stancheva, 1990
SARMATIAN	Late	NO14	Aurila notata	
		NO 13		
	Middle	NO 12		Euxinocythere grave odessosensis
	Early	NO 11	Cytheridea hungarica - Aurila mehesi	Euxinocythere turpe
				Cytheridea hungarica - Aurila mehesi

Fig. 6. Central and Eastern Paratethys Sarmatian substages and correlation based on ostracods.

Another possibility is that the first 5 m of sediments overlaying the Badenian ones, with abundant *Ammonia beccarii*, represent the entire interval of Anomalinoidea dividens biozone in the Central Moesian Platform. Taking into account the later statements, we believe that the oldest foraminiferal zone that might be highlighted in the studied area is *Varidentella reussi* Zone, Early Volhynian in age (except the basal part of this substage).

The following identified biozone is *Elphidium reginum* Zone that has a large development in the studied area, being observed in all analyzed wells. The microfaunal assemblages of the *Elphidium reginum* Zone show differences in abundance related to the type of sediments enclosing them. Hence, the siliciclastic sediments contain in general scarce assemblages with only few foraminifers and ostracods, while in the carbonate rocks the foraminifers and ostracods abundance increases.

The most characteristic species present in the late Volhynian *Elphidium reginum* Zone are:

- (i) In siliciclastic rocks, the found foraminiferal assemblages contain *Ammonia beccarii* (Linnaeus), *Elphidium fichtelianum* (d'Orbigny), *Elphidium macellum aculeatum* Silvestri, *Elphidium macellum macellum* (Fichtel & Moll), and *Elphidium reginum* (d'Orbigny);

- (ii) In carbonate rocks, the foraminifer assemblages are represented by *Pseudotriloculina consobrina* (d'Orbigny), *Varidentella reussi* (Bogdanowicz), *Articulina problema* Bogdanowicz, *Cycloforina karrerri ovata* Serova, *Nonion bogdanowiczii* Voloshinova, *Porosonion subgranosus subgranosus* (Egger), *Buliminella elegantissima* (d'Orbigny), *Bolivina moldavica* Didkowski, *Fursenkoina sarmatica* (Venginski), *Fissurina* sp., *Elphidium aculeatum* (d'Orbigny), *Elphidium hauerinum* (d'Orbigny), *Elphidium josephinum* (d'Orbigny), and *Elphidium reginum* (d'Orbigny), accompanied by the ostracods *Cytheroidea sarmatica* Jiricek, *Cytheridea hungarica* (Zalanyi), *Euxinocythere diafana* (Stancheva), *Euxinocythere naca* (Mehes), *Callistocythere egregia* (Mehes), *Amnicocythere tenuis* Reuss, *Aurila merita* (Zalanyi), *Aurila mehesi* (Zalanyi), *Loxocorniculum hastatum* (Reuss), *Loxocorniculum schmidi* (Cernajsek), *Loxoconcha porosa* Mehes, *Loxoconcha punctatella* (Reuss), *Xestoleberis elongata* Schneider, and *Xestoleberis fuscata* (Schneider), along with serpulids, mysid statoliths, gastropods, bivalves and *Charophyta* algae.

Some of the aforementioned representative taxa are illustrated in Plates 1-4. The distribution of samples in each well and the occurrence of species are presented in the Chronocharts (Figs. 7-15).




Fig. 7. Chronochart Blejești 1 - a.






Fig. 7. Chronochart Blejești 1 - b and c.




Fig. 8. Chronochart Blejesti 3 - **a** and **b**.




Fig. 8. Chronochart Blejești 3 - c and d.




Fig. 9. Chronochart Blejești 4.




Fig. 10. Chronochart Blejești 5.




Fig. 11. Chronochart Blejești 6.




Fig. 12. Chronochart Blejești 7.




Fig. 13. Chronochart Blejești 8.





Fig. 14. Chronochart Blejești 9.






Fig. 15. Chronochart Blejești 10.



4. CONCLUSIONS

Based on the micropaleontological data from 10 drillings located in the south-central part of the W Moesian Platform, we were able to separate the following assemblages:

- (i) Just above the Badenian deposits or overlaying the Cretaceous ones, a transitional zone, lacking foraminifera marker to assign a certain biozone, but with very abundant specimens of the *Ammonia beccarii* foraminifer species, was observed. Possibly, this interval corresponds to the *Anomalinoides dividens* Zone defined by Popescu (1985). This assumption is argued by the identification in the aforementioned interval of the studied wells of the NO 11 *Cytheridea hungarica* - *Aurila mehesi* ostracod zone, which starts at the base of the Sarmatian (=early Volhynian), overlapping the time interval of *Anomalinoides dividens* Foraminiferal Zone.
- (ii) The *Varidentella reussi* Zone was observed in the studied deposits based on the presence of index species, along

with other foraminiferal and ostracod taxa present in the identified microfaunal assemblages. This biozone is indicative for the presence of the lower Volhynian depositional interval in the studied area.

- (iii) The upper Volhynian depositional interval has been pointed out based on the identification of *Elphidium reginum* Foraminiferal Zone. This biozone was identified in two types of sediments. In the siliciclastic deposits intercepted by the wells, the assemblages are very scarce, but contain the index foraminifera taxon, while in the carbonate sediments more diversified and abundant assemblages were identified.

ACKNOWLEDGEMENTS

The authors thank to Prof. Dr. Mihai Brânzila and an anonymous reviewer for their useful suggestions and comments. We also thank to OMV Petrom S.A. Company, for providing the core analyzed and presented herein.

REFERENCES

- ANDRUSOV N. (1917). Pontice skijjarus. *Trudy Geol. Com.* **4** (2).
- BECA C., PRODAN D. (1983). Geologia zăcămintelor de hidrocarburi. Editura didactică și pedagogică, București.
- BRÂNZILĂ M., CHIRA C. (2005). Sarmatian boundary in boreholes of the Moldavian Platform. *Acta Palaeontologica Romaniae* **5**: 17-26.
- BRICEAG A., YANCHILINA A., RYAN W.B.F., OAIE G., MELINTE-DOBRIANESCU M.C. (2016a). Late Pleistocene - Holocene paleoenvironmental changes from the Romanian Black Sea shelf inferred by microfaunal and isotope fluctuations. *International Multidisciplinary Scientific Geoconference SGEM*, **1**: 305-312, doi: 10.5593/sgem2016B11.
- BRICEAG A., MELINTE-DOBRIANESCU M.C., ION G., BALAN S. (2016b). Re-occurrence of *Phyllophora* red algal genus on the Romanian Black Sea shelf. *International Multidisciplinary Scientific Geoconference SGEM*, **2**(3): 813-819, doi: 10.5593/sgem2016B32.
- BRICEAG A., MACALEȚ R., MELINTE-DOBRIANESCU M.C. (2018). Sarmatian palaeoenvironment and bioevents in the Dobrogea region (SE Romania). *Geo-Eco-Marina* **24**: 81-91.
- BRICEAG A., YANCHILINA A., RYAN W., STOICA M., MELINTE-DOBRIANESCU M. (2019). Late Pleistocene to Holocene paleoenvironmental changes in the NW Black Sea. *Journal of Quaternary Science*, **34** (2): 87-100.
- CORNÉE J.-J., MOISSETTE P., SAINT-MARTIN J.-P., KÁZMÉR M., TOTH E., GÖRÖG A., DULAI A., MÜLLER P. (2009). Marine carbonate systems in the Sarmatian (Middle Miocene) of the Central Paratethys: the Zsambek Basin of Hungary. *Sedimentology* **56**(6): 1728-1750.
- DUMITRIU S.D., LONGHIN S., DUBICKA Z., MELINTE-DOBRIANESCU M., PARUCH-KULCZYCKA J., IONESCU V. (2017). Foraminiferal, ostracod, and calcareous nannofossil biostratigraphy of the latest Badenian-Sarmatian interval (Middle Miocene, Paratethys) from Poland, Romania and the Republic of Moldova. *Geologica Carpathica*, **68**(5): 419-444.
- DEBENAY J.P., BENETEAU E., ZHANG J., STOUFF V., GESLIN E., REDOIS F., FERNANDEZ GONZALEZ M. (1998). *Ammonia beccarii* and *Ammonia tepida* (foraminifera): Morphofunctional arguments for their distinction. *Marine Micropaleontology*, **34**: 235-244.
- FILIPESCU S., SILYE L. (2008). New Paratethyan biozones of planktonic foraminifera described from the Middle Miocene of the Transylvanian Basin (Romania). *Geologica Carpathica*, **59**: 537-544.
- FILIPESCU S., MICLEA A., GROSS M.J., HARZHAUSER M., ZÁGORŠEK K., JIPA C., 2014. Early Sarmatian paleoenvironments in the easternmost Pannonian Basin (Borod Depression, Romania) revealed by the micropaleontological data. *Geol. Carpat.* **65** (1): 67-81.
- GEBHARDT H., ZORN I., ROETZEL R. (2009). The initial phase of the early Sarmatian (Middle Miocene) transgression. Foraminiferal and ostracod assemblages from an incised valley fill in the Molasse Basin of Lower Austria. *Austrian Journal of Earth Science*, **102**: 100-119.
- GROSS M. (2002). Middle Miocene ostracods from the Vienna Basin (Badenian/Sarmatian, Austria). Dissertation, Institut für Geologie und Paläontologie Karl-Franzens-Universität Graz, 343 p.

- GÖRÖG A. (1992). Sarmatian foraminifera of the Zambek Basin, Hungary. *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae*, **XXIX**: 31-153.
- HANGANU E., GRIGORESCU D. (1981). Asupra ostracofaunei din nisipurile sarmatiene ale Dobrogei de Sud (Sectorul Cobadin – Cotu Văii). *Studii și cercetări de geologie, geofizică și geografie*, **26** (2): 309-313.
- HARZHAUSER M., PILLER W.E. (2004). Integrated stratigraphy of the Sarmatian (Upper Middle Miocene) in the western Central Paratethys. *Stratigraphy*, **1**: 65-86.
- IONESI B. (1968). Stratigrafia depozitelor miocene de platformă dintre Valea Siretului și Valea Moldovei. Editura Academiei RSR, 396 pp.
- IONESI B., GUEVARA I. (1993). Studiul depozitelor sarmatiene din forajul 1002 Bădeuți (NV Platformei Moldovenești). *Geological Bulletin of Romanian Society*, **14**: 79-87.
- IONESI B., IONESI L. (1968). Contributions to the knowledge of the Ionița C., Ghiță D., Limborea L., Șindilar V., 2016. The first mention of Badenian in Cartojani (Central-Southern area of the Moesian Platform) based on biostratigraphical analyses (microfauna and nannoplankton). AAPG European Regional Conference & Exhibition, Bucharest, May 2016. Abstract Book, 56 p.
- JIPA D.C., OLARIU C. (2009). Dacian Basin: Depositional Architecture and Sedimentary History of a Paratethys Sea. *Geo-Eco-Marina Special Edition* **3**, 268 p.
- JIPA D.C., OLARIU C. (2009). Sediment routing in a semi-enclosed epicontinental sea: Dacian Basin, Paratethys Domain, Late Neogene, Romania. *Global and Planetary Change*, **103**: 193-206.
- JIRÍČEK R. (1985). Die Ostracoden des Pannonien. In: Chronostratigraphie und Neostatotypen Miozän der Zentralen Paratethys. Bd. VII. M6. Pannonien (Slavonien und Serbien): 378-425.
- JIRÍČEK R., ŘÍHA J. (1991). Correlation of Ostracod Zones in the Paratethys and Tethys. *Saito Ho-on Kai Special Publications (Proceedings of Shallow Tethys)* **3**: 435-457.
- KOUBOVÁ I., HUDÁČKOVÁ N. (2010). Foraminiferal successions in the shallow water Sarmatian sediments from the MZ 93 borehole (Vienna Basin, Slovak part). *Acta Geologica Slovaca, Ročník* **2**(1): 47-58.
- MARINESCU F. (1978). Stratigrafia Neogenului superior din sectorul vestic al Bazinului Dacic. Ed. Acad RSR, București, 155 p.
- MELINTE-DOBRIANESCU M., ION G. (2013). Emiliana huxleyi fluctuations and associated microalgae in superficial sediments of the Romanian Black Sea shelf. *Geo-Eco-Marina* **19**: 129-135.
- MURRAY J.W. (1991). Ecology and paleoecology of benthic foraminifera. New York, John Wiley & Sons, 397 p.
- OLTEANU R., JIPA D. (2006). Dacian basin environmental evolution during Upper Neogene within the Paratethys domain. *Geo-Eco-Marina* **12**: 91-105.
- PAPP A., MARINESCU F., SENES J. (1974). M5 - Sarmatien (sensu E. SUESS, 1866). Die Sarmatische Schichten gruppe und ihr Statotypus. Chronostratigraphie und Neostatotypen. Miozän der Zentralen Paratethys. Slowak Akad Wissensch Bratislava, 4.
- PILLER W.E., HARZHAUSER M., MANDIC O. (2007). Miocene Central Paratethys stratigraphy – current status and future directions. *Stratigraphy*, **4**: 151-168.
- POPESCU G. (1995). Contribution to the knowledge of the Sarmatian foraminifera of Romania. *Romanian Journal of Paleontology*, **76**: 85-98.
- POPOV N., KOJUMDIEVA E. (1987). The Miocene in Northeastern Bulgaria (Lithostratigraphic subdivision and geological evolution) [in Bulgarian with English abstract]. *Rev. Bulg. Geol. Soc.* **48** (3): 15-33.
- POPOV S.V., RÖGL F., ROZANOV A.Y., STEININGER F.F., SHCHERBAI. G., KOVÁČ M. (2004). Lithological–Paleogeographic maps of Paratethys. 10 Maps Late Eocene to Pliocene. *Cour Forsch Senck* **250**: 1-46.
- PARASCHIV D. (1979). Platforma Moesică și zăcămintele ei de hidrocarburi. Editura Academiei RSR București, 195 p.
- REINHARDT E.G., PATTERSON R.T., SCHRÖDER ADAMS C.J. (1994). Geoarchaeology of the ancient harbor site of Caesarea Maritima, Israel: Evidence from sedimentology and paleoecology of benthic foraminifera. *Journal of Foraminiferal Research* **24** (1): 37-48.
- ROYDEN L., HORVÁTH F., BURCHFIELD B.C. (1982). Transform faulting, extension and subduction in the Carpathian Pannonian region. *Geological Society of America Bulletin*, **93**: 717-725.
- SAINT MARTIN, J.P., PESTREA, S. (1999). Les constructions à serpules et microbialites du Sarmatien de Moldavie. *Acta Palaeontol. Rom.*, **2**: 463-469.
- SÂNDULESCU M. (1984). Geotectonica României. Editura Tehnică București.
- SCHMID S.M., BERNOULLI D., FÜGENSCHUH B., MATENCO L., SCHEFER S., SCHUSTER R., TISCHLER M., USTASZEWSKI R. (2008). The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units. *Swiss J. Geosci.*, **101**: 139-183.
- SHORT F.T., NECKLES H.A. (1999). The effects of global climate change on sea grasses. *Aquatic Botany* **63** (3-4): 169-196.
- STEININGER F.F., WESSELY G. (2000). From the Tethyan Ocean to the Paratethys Sea: Oligocene to Neogene stratigraphy, paleogeography and paleobiogeography of the circum-Mediterranean region and the Oligocene to Neogene Basin evolution in Austria. *Mitt. Österr. Geol. Ges.* **92**: 95-116.
- TER BORGH M., STOICA M., DONSELLAR M., MATENCO L., KRUGSMAN W. (2014). Miocene connectivity between the Central and Eastern Paratethys: Constraints from the western Dacian Basin. *Palaeogeogr., Palaeoclim., Palaeoecol.* **412**: 45-67.
- ZELINKA J. (1990). A review of the Sarmatian Ostracoda of the Vienna Basin. In: Whatley R., Maybury C. (Eds.): Ostracoda and global events. Chapman & Hall: 263-270.